

Insects >> Butterflies >> Nymphalinae >> Rohana parisatis pseudosiamensis

Rohana parisatis pseudosiamensis



Male underside

Nguy�n-Phung 1985 - The Siamese Black Prince

Koh Samui only.

This species shows extreme sexual dimorphism. The male upperside is without any markings except for a tiny white apical spot. It is not uncommon at forest edges and glades.

Flight Period

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Х	Х	Х	Х	Х	х	х	Х	х	Х	Х	Х

Wingspan

50 - 53mm

Early Stages


Only partially known. More detailed life history information is available from the following resources: S. Igarishi - Life Histories of Asian Butterflies, Volume.2, Page 511

Larval Foodplant

Impatiens violaeflora (Balsaminaceae). Outside Thailand, it has been recorded feeding on various *Celtis* species(Ulmaceae)

Range

Thailand and Malay Peninsular. Other subspecies from Sri Lanka and India to China and Sundaland, including Palawan.



<< Back to Nymphalinae overview

<u>Home</u> | <u>About</u> | <u>Insects</u> | <u>Cercopoidea</u> | <u>Expeditions</u> | <u>Galleries</u> | <u>Contact</u> | <u>Links and Resources</u> | <u>Disclaimer</u> | <u>Privacy</u> | <u>Sitemap</u>

© 2000-2022 Samuibutterflies.com and Les Day Design and Development by tomjansen.net