

SAMUI BUTTERFLIES

www.samuibutterflies.com

by Les Day

[home](#)

[about](#)

[contact](#)

[insects](#) ▼

[expeditions](#) ▼

[galleries](#) ▼

[Insects](#) >> [Butterflies](#) >> [Nymphalinae](#) >> [Cyrestis cocles earli](#)

Cyrestis cocles earli



f. formosa Upperside



f. formosa Underside

Distant 1883 - The Marbled Map

Koh Samui only.

It is primarily a forest species.

This species can be found in two forms, *f. earli* (Distant) and *f. formosa* (C & R Felder) throughout the year, but rarely in great numbers.

Flight Period

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
x	x	x	x	x	x	x	x	x	x	x	x

Wingspan

35 - 50mm

Early Stages

Only partially known. More detailed life history information is available from the following resources: S. Igarishi - Life Histories of Asian Butterflies, Volume.2 , Page 459.

Larval Foodplant

Unidentified species of Moraceae.

Range

N. E. India to S. China, down through the Malay Peninsular, Sumatra and Borneo.



f. earli Upperside



f. earli Underside

[<< Back to Nymphalinae overview](#)