

SAMUI BUTTERFLIES

www.samuibutterflies.com

by Les Day

[home](#)

[about](#)

[contact](#)

[insects](#) ▼

[expeditions](#) ▼

[galleries](#) ▼

[Insects](#) >> [Butterflies](#) >> [Danainae](#) >> [Euploea crameri bremeri](#)

Euploea crameri bremeri



Male upperside



Male upperside

Lucas 1853 - The Spotted Black Crow

Koh Samui only.

A not uncommon species. It can be seen in gardens, feeding on lantana flowers, and in secondary growth.

It can be distinguished by having the apical and inner submarginal spots much enlarged.

There is also a subspecies of *Euploea core*, *E. c. graminifera* which is very similar, but with this species, the males have a small black band on the upperside forewing.

Flight Period

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
		x	x	x	x	x	x	x		x	

Wingspan

70 - 85mm

Early Stages

Known. More detailed life history information is available from the following resource: [ButterflyCircle Blog](#) (courtesy of Dr. Horace Tan.)

Larval Foodplant

Parsonia helicandra (Asclepidaceae)

Range

Burma, through the Peninsular to Sundaland

[<< Back to Danainae overview](#)