

SAMUI BUTTERFLIES

www.samuibutterflies.com

by Les Day

[home](#)

[about](#)

[contact](#)

[insects](#) ▼

[expeditions](#) ▼

[galleries](#) ▼

[Insects](#) >> [Butterflies](#) >> [Nymphalinae](#) >> *Pantoporia hordonia hordonia*

Pantoporia hordonia hordonia



Male upperside



Male underside

[<< Back to Nymphalinae overview](#)

Stoll 1790 - The Common Lascar

Koh Samui only.

Sexes similar.

This species is very similar to the members of *Lasippa* on the upperside. It is to be found at forest edges, and is uncommon. It can be identified from other members of the genus on Samui by having, on the upperside forewing, the inner submarginal line coloured gray, and the outer one, orange.

Flight Period

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
x		x	x	x	x	x		x			x

Wingspan

40 - 50mm

Early Stages

Fully known. More detailed life history information is available from the following resource: [ButterflyCircle Blog](#) (courtesy of Dr. Horace Tan.)

Larval Foodplant

Archidendron clypearia (Mimosaceae)

Range

N. E. India to Indo-China and throughout the Malay Peninsular. Other subspecies from Sri Lanka, Indian peninsular, S. China and Taiwan, also Lesser Sunda Islands.