

SAMUI BUTTERFLIES

www.samuibutterflies.com

by Les Day

[home](#)

[about](#)

[contact](#)

[insects](#) ▼

[expeditions](#) ▼

[galleries](#) ▼

[Insects](#) >> [Butterflies](#) >> [Lycaenidae](#) >> [Jamides bochus nabonassar](#)

Jamides bochus nabonassar



[<< Back to Lycaenidae overview](#)

Fruhstorfer 1916 - Dark Caerulean

This is the only member of the *bochus* subgroup found in Thailand.

It is a small species with strong dark metallic blue uppersides in the male, and is a fast flyer.

It is usually to be found in open country amongst secondary vegetation.

Flight Period

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
x	x	x	x			x	x			x	x

Wingspan

23 - 30mm

Early Stages

Known. More detailed life history information is available from the following resource: [ButterflyCircle Blog](#) (courtesy of Dr. Horace Tan.)

Larval Foodplant

Xylia xylocarpua (Mimosaceae), *Pueraria colletti*, *Pueria mirifica* and *Butea frondosa* (all Papilionaceae)

Range

Peninsular Thailand, Malaysia, Sumatra and Borneo. Other [subspecies](#) can be found from Sri Lanka all the way to the Fiji Islands.