

SAMUI BUTTERFLIES

www.samuibutterflies.com

by Les Day

home

about

contact

insects ▾

expeditions ▾

galleries ▾

Insects >> Butterflies >> Lycaenidae >> Hypolycaena erylus teatus

Hypolycaena erylus teatus



Male upperside



Female upperside

Fruhstorfer 1912 - Common Tit

A common species found throughout the island except in thick forest.

Flight Period

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
x	x	x			x	x	x	x	x	x	x

Wingspan

27 - 32mm

Early Stages

Known. More detailed life history information is available from the following resource: [ButterflyCircle Blog](#) (courtesy of Dr. Horace Tan.)

Larval Foodplant

Senna alata (Caesalpinaceae), *Ceiba pentandra* (Bombacaceae), *Cinnamomum verum*, *Cinnamomum zeylanicum* (both Lauraceae), *Dimocarpus longan* (Sapindaceae), *Sandoricum indicum* (Meliaceae) and *Ziziphus jujuba* (Rhamnaceae)

Range

Peninsular Thailand, Malaysia, Sumatra and Borneo. Many other [subspecies](#) found From Nepal, Sikkim, Myanmar, Vietnam and Northern Thailand, through to New Guinea.



Underside

[<< Back to Lycaenidae overview](#)