

Insects >> Butterflies >> Lycaenidae >> Cigaritis Iohita senema

Cigaritis Iohita senema



Fruhstorfer 1912 - Long-banded Silverline

A common species found in secondary vegetation and fields.

Flight Period

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
X	Х	Х	Х	Х		Х	Х		Х	Х	Х

Wingspan

23 - 28mm

Early Stages

Known. More detailed life history information is available from the following resource: ButterflyCircle Blog (courtesy of Dr. Horace Tan.)

Larval Foodplant

Dioscorea hispida, Dioscoria pentaphylla (both Dioscoreaceae), Psidium guajava (Myrtaceae) and Xylia xylocarpa (Mimosaceae)

Range

Southern Myanmar though to Indo-China, Thailand, down the peninsular through Malaysia to Singapore. Another subspecies can be found from Nepal, through Sikkim, Northern Myanmar to Northern Thailand.



<< Back to **Lycaenidae** overview

Home | About | Insects | Cercopoidea | Expeditions | Galleries | Contact | Links and Resources | Disclaimer | Privacy | Sitemap

© 2000-2022 Samuibutterflies.com and Les Day Design and Development by <u>tomjansen.net</u>