

# SAMUI BUTTERFLIES

www.samuibutterflies.com

by Les Day

[home](#)

[about](#)

[contact](#)

[insects](#) ▼

[expeditions](#) ▼

[galleries](#) ▼

[Insects](#) >> [Butterflies](#) >> [Hesperiidae](#) >> [Hasora malayana](#)

## Hasora malayana



[<< Back to Hesperiidae overview](#)

### C. & R. Felder 1860 - White-Banded Awl

This is probably the most common species of this Genus on Samui. It is a denizen of dense forest.

This species was previously known as *H. taminatus malayana*, but, as its range overlapped that of another subspecies, *H. taminatus bhavara*, which is not allowed under the rules of sympatry, it has now been given specific status.

### Flight Period

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
x	x	x		x		x			x		

### Wingspan

38 - 45mm.

### Early Stages

Fully known. Further details available in Life Histories of Asian Butterflies, volume 2, page 628.

### Larval Foodplant

*Derris scandens* and *Millettia racemosa* (both Papilionaceae)

### Range

S. Myanmar, Peninsular Thailand, West Malaysia, Sumatra, Borneo and Java. Other subspecies found from Sri Lanka, India through to S. China, Taiwan, also Philippines, Moluccas, Lesser Sunda Islands and Waigeu.