Insects >> Butterflies >> Lycaenidae >> Curetis bulis bulis

# **Curetis bulis bulis**



### Westwood 1851 - Bright Sunbeam

This is by far the commoner of the two *Curetis* species found on the Island.

Found in open ground near secondary vegetation. It has broader black tornal markings on the upperside hindwing than *C. santana*.

### **Flight Period**

| Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Х   | Х   | Х   | Х   | Х   | Х   | Х   | Х   | Х   | Х   | Х   | Х   |

### Wingspan

31 - 40mm

## **Early Stages**

Not known

## **Larval Foodplant**

Not known

#### Range

Peninsular Thailand, Malaysia, Sikkim, Assam, Myanmar and Indo-China.



<< Back to **Lycaenidae** overview

<u>Home</u> | <u>About</u> | <u>Insects</u> | <u>Cercopoidea</u> | <u>Expeditions</u> | <u>Galleries</u> | <u>Contact</u> | <u>Links and Resources</u> | <u>Disclaimer</u> | <u>Privacy</u> | <u>Sitemap</u>

© 2000-2022 Samuibutterflies.com and Les Day Design and Development by tomjansen.net