

Insects >> Butterflies >> Lycaenidae >> Catochrysops strabo strabo

# Catochrysops strabo strabo



## Fabricius 1793 - The Forget-me-not

Very similar to C. panormus, but the underside forerwing has a costal spot midway between the cell-end spot and the postdiscal band. It is also found in secondary vegetation.

## **Flight Period**

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
ı	Х	Х	Х				Х	Х	Х	Х	Х	Х

### Wingspan

24 - 30mm

### **Early Stages**

Not fully known.

### **Larval Foodplant**

Cajanus cajan, Cylista scariosa, Desmodium heterocarpon, Phaseolus spp., Phyllodium longipes (all Papilionaceae) and Schleichera oleosa (Sapindaceae)

#### Range

From Ceylon, throughout Indo-China to Taiwan.



<< Back to **Lycaenidae** overview

<u>Home</u> | <u>About</u> | <u>Insects</u> | <u>Cercopoidea</u> | <u>Expeditions</u> | <u>Galleries</u> | <u>Contact</u> | <u>Links and Resources</u> | <u>Disclaimer</u> | <u>Privacy</u> | <u>Sitemap</u>

© 2000-2022 Samuibutterflies.com and Les Day Design and Development by tomjansen.net