

# SAMUIBUTTERFLIES

www.samuibutterflies.com

by Les Day

[home](#)

[about](#)

[contact](#)

[insects](#) ▼

[expeditions](#) ▼

[galleries](#) ▼

[Insects](#) >> [Butterflies](#) >> [Lycaenidae](#) >> [Arhopala centaurus nakula](#)

## Arhopala centaurus nakula



Female upperside



Underside

### C & R Felder 1860 - Common Oakblue

A large species, found primarily in secondary vegetation, near wooded areas. It is the most commonly seen member of the Genus.

### Flight Period

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
x	x	x	x	x	x	x	x	x	x	x	x

### Wingspan

40 - 51mm

### Early Stages

Known. More detailed life history information is available from the following resource: [ButterflyCircle Blog](#) (courtesy of Dr. Horace Tan.)

### Larval Foodplant

*Syzygium fruticosum*, (Myrtaceae), *Mangifera indica* (Anacardiaceae), *Terminalia alata* (Combretaceae), *Quercus ramsbottomii*, *Quercus lamellosa* (Fagaceae) and *Xylia xylocarpa* (Mimosaceae)

### Range

Thailand, Indo-China, Burma, Malaysia, Sumatra and Borneo.

[<< Back to Lycaenidae overview](#)