

SAMUI BUTTERFLIES

www.samuibutterflies.com

by Les Day

[home](#)

[about](#)

[contact](#)

[insects](#) ▼

[expeditions](#) ▼

[galleries](#) ▼

[Insects](#) >> [Butterflies](#) >> [Lycaenidae](#) >> [Rapala pheretima sequeira](#)

Rapala pheretima sequeira



[<< Back to Lycaenidae overview](#)

Distant 1885 - The Copper Flash

Probably the most common, and largest of the Rapala species found on the Islands.

It is a species found at the edge of woodland, in secondary vegetation and in orchards.

Flight Period

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
x	x	x	x	x	x	x	x	x	x	x	x

Wingspan

31 - 35mm

Early Stages

Fully known. More detailed life history information is available from the following resource: [ButterflyCircle Blog](#) (courtesy of Dr. Horace Tan.)

Larval Foodplant

Dimocarpus longan, *Lepisanthes rubiginosa*, *Litchi chinensis*, *Nephelium lappaceum* (all Sapindaceae), *Syzygium fruticosum* (Myrtaceae), *Mangifera indica* (Anacardiaceae) and *Vigna sinensis* (Papilionaceae)

Range

Peninsular Thailand, Malaysia and Singapore. Other subspecies found from India, Nepal through Myanmar to Mainland Thailand.